

Bonemarker normal values for children

Bone marker	Cat. No.	Sex	Unit	Age group					References	
BAP (serum) bone specified alkaline phosphatase	8012	m	U/l ± SD		104 ± 21 0–2 y.	94 ± 17 3–9 y.	114 ± 21 10–14 y.		59 ± 19 15–18 y.	[1]/[2]
		f	U/l ± SD		122 ± 19 0–2 y.	106 ± 16 3–8 y.	112 ± 16 9–12 y.		39 ± 23 13–17 y.	
		Tanner Stage		Tanner I	Tanner II–III	Tanner IV–V				
		m	U/l ± SD	95 ± 22	114 ± 35	121 ± 37				[3]
		f	U/l ± SD	84 ± 23	113 ± 43	79 ± 46				
CICP/PICP (serum) c-terminal propeptid of type I collagen	8003	m/f	ng/ml			295–365 4.0–12.9y.	444 ± 192 13 y.	449 ± 123 14 y.	347 ± 145 15–18 y.	[4]
		m	ng/ml		211–1.310 0–2 y.	147–830 3–9 y.	123–618 10–14 y.	127–189 15–19 y.	77–183 20–50 y.	[1]
		f	ng/ml		230–1.104 0–2 y.	155–568 3–8 y.	122–373 9–12 y.	74–167 13–17 y.	37–133 20–50 y.	
		m/f	ng/ml			113–426 4–13 y.		110–443 14–18 y.		[5]
NTX (serum) N-terminal telopeptide of type I collagen	504836	Tanner Stage		Tanner I	Tanner II–IV	Tanner V				
		m	BCE nmol/l	50.6 (37.2–88.9)	71.7 (32.7–126.0)	18.0 (11.3–50.4)				[6]
		f	BCE nmol/l	53.6 (31.2–90.9)	50.2 (12.9–79.8)	12.8 (9.6–22.2)				
NTX (urine) N-terminal telopeptide of type I collagen	504837			0–1 y.	2–5 y.	6–10 y.	11–15 y.	16–20 y.		
		m/f	BCE nmol/l Creatinin	1639 (102-4769)	689 (34-1752)	497 (90-1356)	429 (34-2158)	192 (34-780)		
Deoxypyridinolin (urine) free DPD	8007	Tanner Stage		Tanner I	Tanner II–III	Tanner IV–V				
		m	nmol/mmol Creatinin	23.5 ± 1.9	25.6 ± 4.0	31.1 ± 9.5				[9]
		f	nmol/mmol Creatinin	21.0 ± 2.1	33.0 ± 5.1	23.2 ± 7.0				
		m/f	nmol/mmol Creatinin	22.5 ± 1.4	28.7 ± 3.2	28.0 ± 6.2				
Helicale peptide (urine) Helicale peptide 620 - 633 from the alpha-I-chain of type I collagen	8022	m/f	ug/mmol Creatinin			804–11305 4–12.9 y.				[5]
Pyridinolin (urine) free Pyd & DPD	8010	m/f	nmol/mmol Creatinin			45.7–85.1 4–12.9 y.				[10]
		m/f	nmol/mmol Creatinin			55.5 ± 6.5 4–12.9 y.	40.8 ± 6.4 13–18 y.			[11]
Pyridinolin (serum)	8019	m/f	nmol/l			2.49 ± 0.12 4–12.9 y.	1.6 ± 1.0 13–18 y.			[11]

Bone marker	Cat. No.	Sex	Unit	Age group						References
				Pre-Pubertal	Early Adolescence	Late Adolescence				
TRAP5b (Serum) Tartrate-Resistant Acid Phosphatase active isoform 5b	8036	status								
		f	U/l	8.1±3.8 1-9 y.	10.0±2.7 10-13 y.	2.3±0.7 14-17 y.				[12]
		m	U/l	6.6±3.6 1-10 y.	9.9±3.3 11-13 y.	3.4±1.4 14-17 y.				[12]

References:

- [1] Tsai et al
Bone Alkaline Phosphatase Isoenzyme and Carboxy-Terminal Propeptide of Type-I Procollagen in Healthy Chinese Girls and Boys.
Clinical Chemistry 45, No. 1, 1999
- [2] Elmlinger MW et al
Significance of bone specific alkaline phosphatase and procollagen-I-peptide as diagnostic markers of bone formation for monitoring growth hormone therapy
at the Congress on Calcium Regulating Hormones and Bone Metabolism from the German Society for Endocrinology, Giessen, Germany, September 29-30, 1995
- [3] Siu L. Hui et al
Difference in Bone Mass between Black and White American Children: Attributable to Body, Build, Sex Hormone Levels, or Bone Turnover?
Journal of Clinical Endocrinology & Metabolism 88 (2), 642 - 649, 2003
- [4] Winterbottom et al
An assay for the C-term. Propeptide of type I collagen.
The Endocrine Society, Anaheim, CA, June 15 - 18, 1994.
- [5] **Quidel In house study**
- [6] Van der Sluis
A Cross-Sectional Study on Biochemical Parameters of bone Turnover and Vitamin D Metabolites in Healthy Dutch Children and Young Adults
Hormon Research 2002;57:170-179.
- [7] F. Rauch
Urinary immunoreactive dpd in children and adolescents: variations with age, sex, and growth velocity.
Scan J. Clin Lab Invest 1996;56: 715 - 719 (Metra Ref # 1540)
- [8] A. Bourdeau
Age Dependent Variations in Healthy Children of Urinary Excretion of Pyridinium Crosslinks.
XII International Conf. On Calcium Regulating Hormones, Australia, Feb. 14-19, 1995.
- [9] A. Conti
Urinary free deoxypyridinoline levels during childhood.
J. Endo. Invest. 21: 318 - 322, 1998. Ref#2574
- [10] Husain, et al
Urinary excretion of pyridinium crosslinks in healthy 4 - 10 year olds.
Arch. Dis. Child 1999, 80: 370 - 373
- [11] A. Colwell
The Renal Clearance of Free and Conjugated Pyridinium Crosslinks of Collagen.
J. of Bone and Min. Res. Vol. 11, Number 12, 1996
- [12] Price HE, Langman CB and Brooks ER.
TRAP5b – profiles in children with chronic kidney disease.
Poster presented at ASBMR (2007).