

Certificate of Analysis

Analysezertifikat

Certificat d'Analyse

Product name	C4d, Human, clone 7H4		
Catalog number	TE2230-100UG		
Lot number	xxxxxXxxxx	Expiry date	MMM YYYY
Volume	1 ml	Amount	100 µg
Formulation	0.2 µm filtered in PBS+0.1%BSA+0.02%NaN3	Concentration	100 µg/ml
Host Species	Mouse IgG1	Conjugate	None
Endotoxin	N.A.	Purification	Protein G
Storage	4°C		

Application notes

	IHC-F	IHC-P	IF	FC	FS	IA	IP	W
Reference #								
Yes	•					•		•
No								
N.D.		•	•	•	•		•	

N.D.= Not Determined; IHC = Immuno histochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.

General Information

Description	The monoclonal antibody 7H4 (also known as clone M4d2) recognizes an epitope on the alpha 2 chain of human C4d. Complement factor C4 (MW 41 kDa), formerly known as Gg protein, consists of an alpha-, beta- and gamma-chain. The classical pathway of complement and the mannose binding lectin (MBL) activation pathway converge at C4. Activated C1, MASP-1 and MASP-2 cleave C4 resulting in the formation of C4a and C4b. The latter can be cleaved by factor I resulting in C4c and C4d, in which step all functional sites are lost. The C4d activation fragment of C4 is an excellent marker for classical complement pathway and MBL pathway activation. The thioester formed between the side chains of Cys1010 and Gln1013 within the C4d region of the α-chain mediates covalent attachment to the target surface bearing activated forms of C1s or MASP. Furthermore, C4d is highly homologous to C3d with over 35% shared amino acid sequence. In a number of diseases such as rheumatoid arthritis (RA), hereditary angioedema (HAE), systemic lupus erythematosus (SLE) and chronic urticaria with hypercomplementemia levels of C4d are significantly elevated in serum or plasma. C4d levels may also be elevated in plasma from patients with a variety of humoral autoimmune diseases in which complement activation is known to occur. Deposition of C4d in peritubular capillaries has been shown to be a sensitive marker for antibody-mediated (humoral) rejection in renal transplant biopsies. The monoclonal antibody 7H4 recognizes C4, C4b and C4d. Antibody 7H4 cross reacts with guinea pig.
Cross reactivity	Guinea pig: Yes.
References	<ol style="list-style-type: none"> Zwirner, J et al; Deposition of complement activation products on plastic-adsorbed immunoglobulins. J Immunol Methods 1989, 124: 121. Zwirner, J et al; Classical pathway of complement activation in normal and diseased human glomeruli. Kidney Int 1989, 36: 1069. Zwirner, J et al; Classical pathway of complement activation in mammalian kidneys. Immunology 1993, 80: 162. Feucht, H et al; Capillary deposition of C4d complement fragment and early renal graft loss. Kidney Int 1993, 43: 1333.
Storage&stability	Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year.

Precautions For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent infringements that might result from the use or derivation of this product.

We hereby certify that the above-stated information is correct and that this product has been successfully tested by the Quality Control Department. This product was released for sale according to the existing specifications. This document has been produced electronically and is valid without a signature.

Approved by Manager of QC

Date

Do you have any questions or comments regarding this product? Please contact us via info@tecomedical.com

www.tecomedical.com



A EURO BIO SCIENTIFIC COMPANY

Switzerland
TECO medical AG
Gewerbstrasse 10
4450 Sissach
Phone +41 61 985 81 00
Fax +41 61 985 81 09
Mail info@tecomedical.com

Germany
TECO medical GmbH
Wasserbreite 57
32257 Bünde
Phone +49 52 23 985 99 99
Fax +49 52 23 985 99 98
Mail info@tecomedical.com

Benelux
TECO medical Benelux BV
Computerweg 22
3542 DR Utrecht, The Netherlands
Phone +31 30 307 87 30
Fax +31 30 307 49 39
Mail benelux@tecomedical.com

Austria
TECO medical AG
Phone 0800 20 40 66
Fax 0800 20 40 55
Mail info@tecomedical.com